

**Quality Sawah Development is the base for Rice Green Revolution in West Africa  
(for IVRDP coordinating team at Kumasi),  
Sustainable intensification and diversification strategies for African rice-based  
cropping systems (for African Rice Congress, Dar es Salaam, 31 July-4 August, 06)  
T. Wakatsuki, Kinki University, Japan**

Guinea, Aug.03

**No Sawah, No Green Revolution**



Inland valley, Sierra Leone, Jan.89

**Water control through Sawah system is prerequisite for Green Revolution in SSA**




Nupe's indigenous rudimentary Sawah system,  
Nigeria, Sep.05



**Fish & Rice**


**CRI/JICA Sawah project, Ghana, Aug.01**

## Production trend (kg/person) of diversified African crops in comparison with Asia



Production  
(kg/person)

Asia



African crops are diverse, even production potential of rice is higher than demand, rice is importing. Wheat has not enough production potential in majority of SSA countries. Rice is also the highest quality cereals in terms of egg protein equivalent among the other 6 crops



Fulbe (maybe also Masai?) cows are not integrated well in the rice farming: Diversity but not good integration in majority of African Agriculture



Nupe farmers' traditional water control systems: Irrigated but rudimentary sawah system because of no availability of animal traction ( and small machinery)





After rice Nupe farmers grow various crops



Sustainable intensive mixed cropping



World record  
Igbo, Nigeria

Sustainable Diversification is not a major problem in SSA. Current major problem is how to realized sustainable intensification: Green Revolution

African nature of Diversity Agriculture may contribute tropical Asian and American agriculture in future.

NERICA rice also may contribute to help Asian rice in future, because of its potential genetic diversity



From plateau to Bangkok plain, July 06

Lagos Airport, Aug. 05

What is Sawah?


Farmers fields are demarcated

No clear demarcation: Land is not injured

Northeastern plateau area in Thailand, July 06

Dar es Salaam airport, July 06


# Farmers' Fields: Diverse and mixed up environments



**mixed up varieties**  
**A B C D E .....**

**Fertilizer, Irrigation, and  
HYV are not effective:  
No Green Revolution**

**Sawah based eco-technology:**  
Diverse but well characterized,  
classified, and improved rice  
environment, **especially for  
water control**




**pure variety A**  
**pure variety B**  
**pure variety C**  
**pure variety D**  
**pure variety E**

**Fig. 7. Successful Integrated Genetic and Natural  
Resource Management, i.e., Agronomy, needs  
classified demarcated land, eco-technologically**


# Biotechnology and Ecotechnology

## Rice variety



Varieties could solve the main problems in Asia  
Is this also true in SSA?

## and Rice with Sawah Systems



Sawah is a man-made, improved rice-growing environment with demarcated, banded, leveled, and puddled fields, **for water control. Sawah is soil based eco-technology**

Because of diverse soil, geology, topography, hydrology, climate, vegetation and socio-cultural conditions, the technologies for sawah development and management are very diverse. Therefore we have to research and develop the technology to accommodate in diverse SSA ecology.

**Fig.2 Rice (variety) and environment (Sawah) improvement Both Bio & Eco-technologies must be developed in balance**



Sawah: Lacking the concept, term and eco-technology. This makes disturbing the balanced approach for rice development in West Africa and SSA last 30 years

## **Confusion in paddy, irrigation, water control, and sawah systems**

Farmers' job

Government Job

- Sawah Hypothesis (1): Antecedent for Green Revolution :  
Are Farmers' field conditions ready to accept irrigation water, fertilizer, and HYV or not?
- Sawah Hypothesis (2): We have to overcome scarce nutrient and water: Sustainable rice productivities under  
Sawah is 10-15 times higher than upland rice fields
- Must remember that lacking the concept & term, "Tsunami" made the Sumatra disaster enormous

No proper English/French ecotechnological concept and term to improve farmers' rice fields,  
Sawah or SUIDEN (in Japanese)

Suiden (Japanese) = **SAWAH** (Malay-Indonesian)


	English	Indonesian	Chinese (漢字)
Plant	Rice	Nasi	米, 飯, 稻
Biotechnology	Paddy	Padi	稻, 粳
Environment	(Paddy) ?	Sawah	水田
Ecotechnology			



Weeds are stronger: upland rice, Bida



No ecotechnology measures



Inland Valley, Sierra Leone

Nupe's traditional partial water control system



Nupe's indigenous partial water control system



Once Sawah systems are developed by farmers' self-support efforts and water is controlled, majority of HYV can produce higher than 5 t/ha

Table Mean gain yield of 23 rice cultivars in low land ecologies at **low (LIL)** and **high input levels (HIL)**, Ashanti, Ghana (Ofori & Wakatsuki, 2005)


Entry No. Cultivar			← ECOTECHNOLOGICAL YIELD IMPROVEMENT					
			<u>Irrigated Sawah</u>		<u>Rainfed sawah</u>		<u>Upland like fields</u>	
			HIL	LIL	HIL	LIL	HIL	LIL
			(t/ha)		(t/ha)		(t/ha)	
BIOTECHNOLOGICAL IMPROVEMENT	1	WAB	4.6	2.9	2.8	1.6	2.1	0.6
	2	EMOK	4.0	2.8	2.9	1.3	1.4	0.5
	3	PSBRC34	7.7	3.5	3.0	2.1	2.0	0.4
	4	PSBRC54	8.0	3.7	3.8	2.1	1.7	0.4
	5	PSBRC66	5.7	3.3	3.8	2.0	1.8	0.4
	6	BOAK189	7.0	3.8	3.7	2.0	1.4	0.3
	7	WITA 8	7.8	4.2	4.4	2.1	1.8	0.5
	8	Tox3108	7.1	4.1	4.0	2.3	2.3	0.6
	9	IR5558	7.9	4.0	3.8	2.0	1.8	0.5
	10	IR58088	7.7	4.0	3.7	1.8	1.4	0.3
	11	IR54742	7.7	4.3	4.0	2.2	1.9	0.4
	12	C123CU	6.9	4.1	4.2	1.9	2.0	0.4
	13	CT9737	6.5	4.0	4.0	1.7	1.9	0.6
	14	CT8003	7.3	3.8	3.8	1.7	2.0	0.5
	15	CT9737-P	8.2	4.0	4.3	1.8	1.2	0.5
	16	WITA1	7.6	3.6	3.3	1.8	0.9	0.3
	17	WITA3	7.6	3.5	4.1	2.0	1.3	0.5
	18	WITA4	8.0	4.1	3.7	2.1	1.5	0.3
	19	WITA6	8.0	3.5	4.0	2.3	1.4	0.3
	20	WITA7	7.3	3.7	3.8	2.2	2.0	0.4
	21	WITA9	7.6	4.4	4.5	2.8	2.0	0.6
	22	WITA12	7.6	4.0	3.8	1.9	1.8	0.4
	23	GK88	7.5	3.8	3.5	2.0	1.8	0.5
Mean (n=23)			7.2	3.8	3.8	2.0	1.7	0.4
Range			(4.0-8.2)	(2.8-4.4)	(2.8-4.5)	(1.3-2.8)	(0.9-2.3)	(0.3-0.6)
SD			1.51	0.81	0.81	0.45	0.44	0.12

Entry 1- 7 : Early - maturing cultivars、Entry 8-23: intermediate - maturing cultivars



## IGNRM for Green Revolution technologies :

Water, Fertilizer and Soil mgt, Legume, Manure, animal traction ,power tiller, fish and rice vegetable after rice, BNF, Iron toxicity, P deficiency, IPM, Striga, Weed mgt, Cover crops, Cropping systems research, CH<sub>4</sub> emission, carbon sequestration, etc. Crop Physiology, Long-term field experiments



**Missing link  
is “sawah”  
in SSAfrica**


High yield  
High quality  
High tolerance

**Ecotechnologically demarcated field for water**

**Management has to be existed:**

High quality leveled rice field ( $\pm 5\text{cm}$ )  
High quality banded rice field (no leaking)  
High quality puddled rice field (nursery)

**Fig. 8. Concept of Integrated Genetic and Natural Resources Management (IGNRM) for green revolution technology :**  
**Missing link is Sawah which is lacking in majority of famers' fields**



West Africa map showing selected sampling sites of lowland soils. (Buri and Wakatsuki, 2000)




Mean values of fertility properties of inland valleys (IVS) and flood plains (FLP) of West Africa in comparison with lowland top-soils of tropical Asia and Japan

Location	Total C (%)	Total N (%)	Available P (ppm)**	Exchangeable Cation (cmol/kg)				Sand (%)	Clay (%)	CEC /Clay
				Ca	K	Mg	eCEC			
IVS	1.3	0.11	9	1.9	0.3	0.9	4.2	60	17	25
FLP	1.1	0.10	7	5.6	0.5	2.7	10.3	48	29	36
T. Asia*	1.4	0.13	18	10.4	0.4	5.5	17.8	34	38	47
Japan	3.3	0.29	57	9.3	0.4	2.8	12.9	49	21	61


\*Kawaguchi and Kyuma (529 sites), 1977,\*\* Bray II.

Source: Hirose and Wakatsuki (268 sites), 1997.


Sulfate - S  
(mg/kg)




Available Zn  
(mg/kg)



Available Cu  
(mg/kg)



Available Ni  
(mg/kg)



■ : River flood plains

■ : Inland valley swamps

S & Zn Deficiency: Frequency distribution of topsoil (0-15cm) available nutrients in West Africa lowlands. (Buri & Wakatsuki. 2001)

# How can we overcome such low level nutrients & scarce water in Sub Sahara West Africa

- To develop lowland sawah is the answer.
- The integrated management of lowland & upland, for example, watershed agro-forestry, is also key eco-technology
- The core region of West Africa has similar climate, soil, hydrology, and crops to northeastern Thailand: The important site in Asian African collaboration in future



Sawah hypothesis (II): Sustainable Productivity of lowland Sawah fields are more than 10 times higher than Upland Fields: This is not experimented results scientifically, but experienced results in Asia



1ha sawah = 10-15ha of upland
-------------------------------

	Upland	Lowland(Sawah)
Area (%)	95 %	5 %
Productivity (t/ha)	1-3 <span style="border: 1px solid red; border-radius: 50%; padding: 2px;"><math>1 \leq</math></span> **	3-6 <span style="border: 1px solid red; border-radius: 50%; padding: 2px;"><math>2^{**}</math></span>
Required area for sustainable 1 ha cropping	5 ha	: 1 ha

\* Assuming 2 years cultivation and 8 years fallow in sustainable upland cultivation, while no fallow in sawah

\*\*In Case of No fertilization

**(1) The optimum landuse pattern and landscape management practices optimize the geological fertilization through the control of optimum hydrology in watershed**




**(2) Sawah systems as multi-functional constructed wetlands**

## Macro- and Micro-scale Ecological Mechanisms of Intensive Sustainability of Lowland Sawah Systems

(1) Geological Fertilization: lowland can receive water, nutrients, and fertile topsoils from uplands.

(2) Multi-functional Constructed Wetlands for control weed and enhanced Supply of N, P, Si, and other Nutrients


**(1) The optimum landuse pattern and landscape management practices optimize the geological fertilization through the control of optimum hydrology in watershed**



**Concept of Watershed Eco-technology, i.e. Watershed Agroforestry: Multi-functional Sawah type wetland is a key component**




Japanese Inland Valley (SATO-YAMA systems): Integration of Forest, Pond and lowland Sawah in watersheds



**Sawah is Multi-Functional Wetland: Rice, Algae, and Microbes' Complex Ecosystems**




# Topsoil, water, and nutrients accumulation through watershed agroforestry



(2) Sawah systems as multi-functional constructed wetlands

# Examples of ecotecnological research & Developemnt



**Two benchmark watersheds in Ghana & Nigeria. Map shows countries with major ethnic groups in West Africa**



# CRI-CSIR/JICA Sawah project for Integrated watershed management, 1997-2001










Although hydrology is the base for success of Sawah, the performance of various ecotechnologies in watershed can be evaluated by water flow.







The conour lines show the depth of ground or surface water level in cm.

Cross-section of topography, rainfall pattern, ground/surface water and land use dynamics in stream flow inland valley (Transect 5), Dwinyan watershed, Ashanti Region, Ghana.



Top-survey, Inland valley, Ashanti, Ghana



Simple barrage by farmers' efforts



Canal construction by farmers



Spring Irrigated Rudimentary Sawah, Nupe





Sawah construction can be done by farmers' self-support efforts



Manual Leveling needs hard-works for  
Sawah system construction



Power-tiller operated leveling, Ashanti, Ghana  
Sawah can/must be produced by farmers



The leveling needs skilled & concerted works





# Sawah is ecotechnology based Multi-Functional constructed Wetland: Production, Environment, and Cultural landscape

Termite mound




Inland valley, Ashanti, Ghana



Rice farmer's field demarcation based on soil, water, and topography are the starting point for scientific observation, technology generation, and application.

Upland leveling & bunding  
are limited only for good soil

Lowland is the  
target for leveling  
and bunding.



Water table and water management continuum(WARDA2004, 2006)

# Can watersheds of in SSA sustain Sawah system?

## High rate of soil erosion and lowland sawah soil formation can be compensated by high rate of soil formation: Again Ecological Balance is a Key





Fig. 1. Rate of soils erosion in the world (Walling1983)



**Irrigation options: Sawah to sawah/contour bund water harvesting, spring, dyke, river, pump, peripheral canal, interceptor canal, tank**

Lowland sawah development priority

$[S] > [L] > [F] > [W] > [U]$

**Concept of Characterization and quantitative mapping of Lowland diversity for sawah development (bunded, leveled, puddled rice land) . depending on the watershed land use, lowland topography, soil, hydrology and Agroecological zones**

**Table 2.** Estimation of rice production trend by each rice ecology in West Africa during 1984-1999/2003. The **2015 estimation is by the author** (WARDA strategic plan in 1988, African rice initiative 2002, Sakurai 2003, WARDA strategic plan 2004, FAOSTAT 2005)

	Area (million ha)			Production (million ton/y)			Yield (t/ha)		
	1984	1999/03	2015	1984	1999/03	2015	1984	1999/03	2015
Upland contribution (%)	1.5 57%	1.8 40%	2.0 30%	1.5 42%	1.8 23%	2.0 13%	1 No yield increase	1	1
Rainfed lowland	0.53	1.8	3.0	0.75	3.4	7.0	1.4	2.0	2.4
Irrigated lowland	0.23	0.56	0.80	0.64	1.9	3.0	2.8	3.4	3.8
Total	2.6	4.7	6.0	3.4	7.7	14	1.3	1.6	2.4



Farmers' sawah fields are the most important infrastructure: farmers' fields come the first

Japanese Experiences




Figure 6. Rice yields & sawah area of historical path in Japan in comparison with rice yields in Asia & Africa

# Distribution of lowlands and potential irrigated sawah in SSA (Hekstra, Andriesse, Windmeijer 1983 & 1993, Irrigated Sawah area estimate by Wakatsuki 2002)

Classification	Area (million ha)		Percentage(%)
Coastal swamps	16.5	(5?)	7
Inland basins	107.5	(4?)	45
Flood plains	30.0	(10?)	12
Inland valleys	85.0	(15?)	36

Possible area of sawah development (million ha)

Max 20million ha (Estimated sawah area came from the relative amount of water cycle in Monsoon Asia, which has 100 million ha of sawah)

# Biotechnology (seed) & Ecotechnology (sawah)

Need Balanced Research and Development


We are going to have many good varieties but farmers fields are/**were** not ready to accept them in SSA

## NEGLECTED PRIORITY MATTERS

- **Massive On The Job farmers' Training program for Sawah based rice technology: Asian African collaboration**
- **Water, soil, and topographic characterization and mapping of Inland Valley Watersheds and flood plain for sustainable lowland sawah development**
- **In Asia, lowland availability is major limiting factor, but it seems water availability in relation to topography and climate will be major limiting factor in SSA's Sawah Development**



# Integrated Genetic & National Resource Management Technology: Need clear concept and target, which can be examined and monitored



**Monitorable Target of Increase**

$$(4-2) \text{ t/ha} \times 4 \times 10^5 + 8 \times 2 \times 10^5 \\ \rightarrow 4.8 \times 10^6 \text{ t/y}$$

# Conclusion: Integrated ecotechnology and biotechnology based *African Green Revolution*



These are still rudimentary Sawah (Bida, Nigeria), but the number of sawah based rice farmers who are consciously developing water & soil management systems are steadily increasing in past 15 years. Prerequisite will be soon satisfied therefore within 10-20 years, the green revolution will be realized in SSA, especially in West Africa, if proper balanced strategy & policy were adopted for African green revolution

# Conclusion

- Sawah is the rice fields to improve water control. The improvement can be done by farmers' efforts
- The farmer based sawah ecotechnology can fill the present big gaps between rainfed rice fields and full irrigated rice fields in West Africa.

The Sawah ecotechnology must be researched by and transferred from researchers to engineers, extension officers, and farmers through intensive and extensive demonstration and on the job training programs



SARI, Tamale, farmers' field training school  
Upland NERICA: 1-1.5t/ha, 16 Aug 06



Rice consumption survey



Garizegu village, near SARI, French assisted contour bund system: 2-3t/ha

# Garizegu village: French assisted contour bund and check dam systems

Yield never over 3t/ha under high yielding varieties and fertilization

